

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

STORM[®] SOFT BAIT
RODENTICIDE

ACTIVE CONSTITUENT: 0.05 g/kg FLOCOUMAFEN

READY TO USE BAIT – KILLS IN A SINGLE FEED

Controls rats and mice in and around industrial, commercial,
agricultural and domestic buildings.
Controls rats and mice resistant to warfarin.

CONTENTS: 5KG

BASF Australia Ltd ABN 62 008 437 867
Level 12, 28 Freshwater Place Southbank VICTORIA 3006

Website: www.agro.com.au

© BASF, 2016[®] = Registered trademark of BASF

APVMA Approval No.: 80663/101357

GENERAL INFORMATION

STORM Soft Bait can be used both indoors and outdoors. Rats and mice that are resistant to warfarin are controlled and no prebaiting is required. A lethal dose is readily consumed in a single feed but death may not occur for up to seven days. DO NOT place bait on or near exposed food or food utensils, food preparation areas or food storage areas.

DIRECTIONS FOR USE

RESTRAINTS

DO NOT apply baits to crops

DO NOT exceed 3 m between bait stations for mice, or 9 m for rats.

DO NOT place baits in the open.

| SITUATION | PEST | RATE | CRITICAL COMMENTS |
|--|---|--|--|
| In and around industrial, commercial, agricultural domestic, and public service buildings, in sewers, animal houses, wharves and around grain terminals. | Rats - <i>Rattus rattus</i> also known as Black rat, Roof, rat, Ship rat or, Fruit rat and <i>Rattus norvegicus</i> also known as the Brown rat or Norway rat | 3 - 5 STORM Soft Baits at each bait location 1 - 2 STORM Soft Baits in each rat hole/burrow | <p>Select suitable bait locations, such as in active rat holes/burrows, along rat runs and under rubbish. Try to establish a barrier of bait locations between living and feeding areas. Secure 3 - 5 STORM Soft Baits at each bait location. STORM Soft Baits should be placed in areas inaccessible to non-target species, such as in bait trays, inside bait boxes or within enclosed spaces.</p> <p>1 - 2 STORM Soft Baits can also be placed within each active rodent hole/burrow.</p> <p>On initial application, an inspection of bait locations after 3 days may be undertaken to determine activity in bait locations and new locations may be selected if necessary. A subsequent inspection may also be undertaken again after 4 days to replace any soft baits which have been eaten.</p> <p>For severe infestations it is recommended that repeat inspections of all bait locations are conducted at 7 day intervals, replacing fresh STORM Soft Baits only when STORM Soft Baits have been eaten and until all signs of rat activity have disappeared.</p> <p>Normally only 3 to 4 baiting rounds (pulses) are required for control.</p> |
| | Mice – <i>Mus musculus</i> also known as the House mouse | 1 - 3 STORM Soft Baits at each bait location | <p>Mice have very limited home ranges and do not need to drink. They are more difficult to control than rats because their feeding patterns are more erratic. Selection of bait locations is therefore even more important than for rats.</p> <p>Secure 1 - 3 STORM Soft Baits in areas inaccessible to non-target species, such as in bait trays, inside bait boxes or within enclosed spaces, spaced 2-3 metres apart. Bait locations should be in areas where activity is obvious, particularly where droppings are seen. Inspect bait locations and replace STORM Soft Baits as for rats until all signs of mouse activity have disappeared.</p> |

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

GENERAL INSTRUCTIONS

Place STORM Soft Baits in areas frequented by the target pests. DO NOT place baits in areas accessible to domestic animals, children, pets and wildlife. DO NOT place baits in areas liable to flooding. Alternative food sources should be removed or reduced as much as possible.

STORM Soft Baits contain Bitrex® (a taste deterrent) to help prevent accidental human and pet consumption.

PRECAUTIONS

DO NOT re-handle product unless wearing chemical resistant gloves.

DO NOT place baits in the open.

DO NOT allow baits to contaminated foodstuffs or food intended for human or animal consumption.

DO NOT remove bait from sachet.

DO NOT use in areas accessible to children.

PROTECTION OF LIVESTOCK, WILDLIFE, FISH, CRUSTACEANS AND OTHERS

This bait is hazardous to pigs, cats, dogs, poultry, birds and wildlife. DO NOT place baits in locations which are accessible to other animals or birds. DO NOT contaminate ponds, streams, rivers or waterways with the bait or its used containers. The product is restricted to use in and around buildings (within 2 metres) or enclosed spaces, e.g., drains. In enclosed roof spaces and areas where possums and other non-target animals frequent, baits should be secured inside proper bait stations, plastic jars or plastic tubes. The use of this product for the control of protected native rodents requires permission from wildlife authorities.

STORAGE AND DISPOSAL

Store in the closed original container in a safe place. DO NOT store for prolonged periods in direct sunlight. Containers which have held bait should not be used for any other purpose. Pest Control Operators should dispose of empty containers, used bait stations and unused or untaken bait by burying in an approved dump. In a domestic situation, minimise the presence of excess baits. Where unused baits are in good condition, return to original container for reuse. For empty containers, damaged or unwanted baits, wrap securely and dispose of in household refuse bin. Look for dead rats or mice and burn or bury them. DO NOT place dead rats or mice in refuse bins or rubbish tips. Ensure baits are removed when rodent activity has ceased.

SAFETY DIRECTIONS

May irritate eyes and skin. Avoid contact with eyes and skin. When using the product, wear elbow-length chemical resistant gloves. Wash hands after use. After each day's use, wash gloves.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Telephone 13 11 26. Vitamin K (Phytomenadione) is antidotal.

NOTE TO DOCTOR

The active constituent flocoumafen is an indirect blood anticoagulant. Ingestion of a relatively large quantity of bait would be required before a toxic effect is expected. In general, an increased tendency to bleed could indicate poisoning. Vitamin K1 (phytomenadione) is antidotal. In cases of suspected poisoning, determine prothrombin time not less than 18 hours after consumption. If elevation has occurred, administer Vitamin K1, intravenously in divided doses, 40 mg for adults, 20 mg for children daily. Treatment may be required daily for several weeks. Withdraw antidote with care and monitor prothrombin times continuously for 4 days or more after such withdrawal.

DOMESTIC ANIMAL POISONING

In cases of accidental poisoning by STORM Soft Bait consult a veterinarian who will advise of suitable Vitamin K1 antidote therapy.

SDS

Additional information is listed in the Safety Data Sheet available from your supplier.

This product does not require a hazard warning label in accordance with GHS

CONDITIONS OF SALE

All conditions and warranties rights and remedies implied by law or arising in contract or tort whether due to the negligence of BASF Australia Ltd ABN 62008437867 or otherwise are hereby expressly excluded so far as the same may legally be done provided however that any rights of the Buyer pursuant to non excludable conditions or warranties of the Competition and Consumer Act 2010 or any relevant legislation of any State are expressly preserved but the liability of BASF Australia Ltd or any intermediate Seller pursuant thereto shall be limited if so permitted by the said legislation to the replacement of the goods sold or the supply of equivalent goods and all liability for indirect or consequential loss or damage of whatsoever nature is expressly excluded. This product must be used or applied strictly in accordance with the instructions appearing hereon. This product is solely sold for use in Australia and must not be exported without the prior written consent of BASF Australia Ltd.

® Storm is a registered trademark of BASF

® Bitrex is a registered trademark of the Macfarlan Smith Limited

Batch No:

Date of Manufacture:

BASF Australia Ltd
ABN 62 008 437 867
Level 12, 28 Freshwater Place
Southbank VICTORIA 3006

FOR SPECIALIST ADVICE IN AN EMERGENCY ONLY PHONE 1800 803 440 TOLL FREE-ALL HOURS-AUSTRALIA WIDE.